

# The Structure of the Atom

**BIG Idea** Atoms are the fundamental building blocks of matter.

## 4.1 Early Ideas About Matter

**MAIN Idea** The ancient Greeks tried to explain matter, but the scientific study of the atom began with John Dalton in the early 1800s.

## 4.2 Defining the Atom

**MAIN Idea** An atom is made of a nucleus containing protons and neutrons; electrons move around the nucleus.

## 4.3 How Atoms Differ

**MAIN Idea** The number of protons and the mass number define the type of atom.

## 4.4 Unstable Nuclei and Radioactive Decay

**MAIN Idea** Unstable atoms emit radiation to gain stability.


## ChemFacts

- Diamond and graphite are both made out of the same element—carbon.
- When graphite was first discovered, it was mistaken for lead. That is why pencils are sometimes called lead pencils.
- There are about  $5 \times 10^{22}$  atoms of carbon in the graphite portion of a pencil.

Graphite surface



Carbon atom



Carbon nucleus



# Start-Up Activities

## LAUNCH Lab

### How can the effects of electric charges be observed?

Electric charge plays an important role in atomic structure.



#### Procedure

1. Read and complete the lab safety form.
2. Cut out small round pieces of **paper** using a **hole punch**, and spread them out on a table.
3. Run a **plastic comb** through your hair. Bring the comb close to the pieces of paper. Record your observations.
4. Obtain two **10-cm pieces of tape**. Fold a 1-cm portion of each piece back on itself to form a handle. Stick both pieces of tape firmly to your desktop. Then, quickly pull both pieces off of the desktop and bring them close together so that their nonsticky sides face each other. Record your observations.
5. Stick a third piece of tape to your desktop. Stick a fourth piece of tape on top of it. Quickly pull the pieces of tape off of the desktop and pull them apart. Bring the two pieces close together so that their nonsticky sides face each other. Record your observations.

#### Analysis


1. **Interpret** your observations using your knowledge of electric charge. Determine which charges are similar and which ones are different.
2. **Explain** how you can tell.
3. **Infer** why neutral pieces of paper were attracted to the charged comb in Step 3 above.

**Inquiry** How can you relate the different charges you have observed to the structure of matter?


## FOLDABLES™ Study Organizer

**The Atom** Make the following Foldable to help you organize your study of the structure of the atom.

- ▶ **STEP 1** Fold a sheet of paper in half lengthwise. Make the back edge about 2 cm longer than the front edge.




- ▶ **Step 2** Fold into thirds.



- ▶ **Step 3** Unfold and cut along one fold line to make one small tab and one large tab.



- ▶ **Step 4** Label as shown.



**FOLDABLES** Use this Foldable with Section 4.2. As you read the section, record information about the atom and its parts.

## Chemistry Online

Visit [glencoe.com](http://glencoe.com) to:

- ▶ study the entire chapter online
- ▶ explore **Concepts in Motion**
- ▶ take Self-Check Quizzes
- ▶ use the Personal Tutor to work Example Problems step-by-step
- ▶ access Web Links for more information, projects, and activities
- ▶ find the Try at Home Lab, Comparing Atom Sizes

## Section 4.1

### Objectives

- ▶ **Compare and contrast** the atomic models of Democritus, Aristotle, and Dalton.
- ▶ **Understand** how Dalton's theory explains the conservation of mass.

### Review Vocabulary

**theory:** an explanation supported by many experiments; is still subject to new experimental data, can be modified, and is considered successful if it can be used to make predictions that are true

### New Vocabulary

Dalton's atomic theory

## Early Ideas About Matter


**MAIN Idea** The ancient Greeks tried to explain matter, but the scientific study of the atom began with John Dalton in the early 1800s.

**Real-World Reading Link** A football team might practice and experiment with different plays in order to develop the best-possible game plan. As they see the results of their plans, coaches can make adjustments to refine the team's play. Similarly, scientists over the last 200 years have experimented with different models of the atom, refining their models as they collected new data.

### Greek Philosophers

Science as we know it today did not exist several thousand years ago. No one knew what a controlled experiment was, and there were few tools for scientific exploration. In this setting, the power of the mind and intellectual thought were considered the primary avenues to the truth. Curiosity sparked the interest of scholarly thinkers known as philosophers who considered the many mysteries of life. As they speculated about the nature of matter, many of the philosophers formulated explanations based on their own life experiences.

Many of them concluded that matter was composed of things such as earth, water, air, and fire, as shown in **Figure 4.1**. It was also commonly accepted that matter could be endlessly divided into smaller and smaller pieces. While these early ideas were creative, there was no method available to test their validity.



■ **Figure 4.1** Many Greek philosophers thought that matter was composed of four elements: earth, air, water, and fire. They also associated properties with each element. The pairing of opposite properties, such as hot and cold, and wet and dry, mirrored the symmetry they observed in nature. These early ideas were incorrect and non-scientific.


**Democritus** The Greek philosopher Democritus (460–370 B.C.) was the first person to propose the idea that matter was not infinitely divisible. He believed matter was made up of tiny individual particles called *atomos*, from which the English word *atom* is derived. Democritus believed that atoms could not be created, destroyed, or further divided. Democritus and a summary of his ideas are shown in **Table 4.1**.

While a number of Democritus’s ideas do not agree with modern atomic theory, his belief in the existence of atoms was amazingly ahead of his time. However, his ideas were met with criticism from other philosophers who asked, “What holds the atoms together?” Democritus could not answer the question.

**Aristotle** Other criticisms came from Aristotle (384–322 B.C.), one of the most influential Greek philosophers. He rejected the notion of atoms because it did not agree with his own ideas about nature. One of Aristotle’s major criticisms concerned the idea that atoms moved through empty space. He did not believe that empty space could exist. His ideas are also presented in **Table 4.1**. Because Aristotle was one of the most influential philosophers of his time, Democritus’s atomic theory was eventually rejected.

In fairness to Democritus, it was impossible for him or anyone else of his time to determine what held the atoms together. More than two thousand years would pass before scientists would know the answer. However, it is important to realize that Democritus’s ideas were just that—ideas, not science. Without the ability to conduct controlled experiments, Democritus could not test the validity of his ideas.

Unfortunately for the advancement of science, Aristotle was able to gain wide acceptance for his ideas on nature—ideas that denied the existence of atoms. Incredibly, the influence of Aristotle was so great and the development of science so primitive that his denial of the existence of atoms went largely unchallenged for two thousand years!

 **Reading Check Infer** why it was hard for Democritus to defend his ideas.

## VOCABULARY



### WORD ORIGIN

#### Atom

comes from the Greek word *atomos*, meaning *indivisible*

**Table 4.1**


## Ancient Greek Ideas About Matter

Philosopher	Ideas
<p><b>Democritus</b> (460–370 B.C.)</p> 	<ul style="list-style-type: none"> <li>• Matter is composed of atoms, which move through empty space.</li> <li>• Atoms are solid, homogeneous, indestructible, and indivisible.</li> <li>• Different kinds of atoms have different sizes and shapes.</li> <li>• Size, shape, and movement of atoms determine the properties of matter.</li> </ul>
<p><b>Aristotle</b> (384–322 B.C.)</p> 	<ul style="list-style-type: none"> <li>• Empty space cannot exist.</li> <li>• Matter is made of earth, fire, air, and water.</li> </ul>



**Table 4.2**

**Dalton's Atomic Theory**


Scientist	Ideas
<p><b>Dalton</b> (1766–1844)</p> 	<ul style="list-style-type: none"> <li>• Matter is composed of extremely small particles called atoms.</li> <li>• Atoms are indivisible and indestructible.</li> <li>• Atoms of a given element are identical in size, mass, and chemical properties.</li> <li>• Atoms of a specific element are different from those of another element.</li> <li>• Different atoms combine in simple whole-number ratios to form compounds.</li> <li>• In a chemical reaction, atoms are separated, combined or rearranged.</li> </ul>


**John Dalton** Although the concept of the atom was revived in the eighteenth century, it took another hundred years before significant progress was made. The work done in the nineteenth century by John Dalton (1766–1844), a schoolteacher in England, marks the beginning of the development of modern atomic theory. Dalton revived and revised Democritus's ideas based on the results of scientific research he conducted. In many ways, Democritus's and Dalton's ideas are similar.

Thanks to advancements in science since Democritus's day, Dalton was able to perform experiments that allowed him to refine and support his hypotheses. He studied numerous chemical reactions, making careful observations and measurements along the way. He was able to determine the mass ratios of the elements involved in those reactions. The results of his research are known as **Dalton's atomic theory**, which he proposed in 1831. The main points of his theory are summarized in **Table 4.2**. Dalton published his ideas in a book, an extract of which is shown in **Figure 4.2**.

 **Reading Check** Compare and contrast Democritus' and Dalton's ideas.

■ **Figure 4.2** In his book *A New System of Chemical Philosophy*, John Dalton presented his symbols for the elements known at that time and their possible combinations.





■ **Figure 4.3** When atoms of two or more elements combine to form a compound, the number of atoms of each element is conserved. Thus, the mass is conserved as well.

**Conservation of mass** Recall from Chapter 3 that the law of conservation of mass states that mass is conserved in any process, such as a chemical reaction. Dalton’s atomic theory easily explains that the conservation of mass in chemical reactions is the result of the separation, combination, or rearrangement of atoms—atoms that are not created, destroyed, or divided in the process. The formation of a compound from the combining of elements and the conservation of mass during the process are shown in **Figure 4.3**. The number of atoms of each type is the same before and after the reaction. Dalton’s convincing experimental evidence and clear explanation of the composition of compounds, and conservation of mass led to the general acceptance of his atomic theory.

Dalton’s atomic theory was a huge step toward the current atomic model of matter. However, not all of Dalton’s theory was accurate. As is often the case in science, Dalton’s theory had to be revised as additional information was learned that could not be explained by the theory. As you will learn in this chapter, Dalton was wrong about atoms being indivisible. Atoms are divisible into several subatomic particles. Dalton was also wrong about all atoms of a given element having identical properties. Atoms of the same element can have slightly different masses.

## Section 4.1 Assessment

### Section Summary

- ▶ Democritus was the first person to propose the existence of atoms.
- ▶ According to Democritus, atoms are solid, homogeneous, and indivisible.
- ▶ Aristotle did not believe in the existence of atoms.
- ▶ John Dalton’s atomic theory is based on numerous scientific experiments.

1. **MAIN** **Contrast** the methods used by the Greek philosophers and Dalton to study the atom.
2. **Define** *atom* using your own words.
3. **Summarize** Dalton’s atomic theory.
4. **Explain** how Dalton’s theory of the atom and the conservation of mass are related.
5. **Apply** Six atoms of Element A combine with 15 atoms of Element B to produce six compound particles. How many atoms of Elements A and B does each particle contain? Are all of the atoms used to form compounds?
6. **Design** a concept map that compares and contrasts the atomic ideas proposed by Democritus and John Dalton.

## Section 4.2

### Objectives

- **Define** atom.
- **Distinguish** between the subatomic particles in terms of relative charge and mass.
- **Describe** the structure of the atom, including the locations of the subatomic particles.

### Review Vocabulary

**model:** a visual, verbal, and/or mathematical explanation of data collected from many experiments

### New Vocabulary

atom  
cathode ray  
electron  
nucleus  
proton  
neutron

## Defining the Atom

**MAIN Idea** An atom is made of a nucleus containing protons and neutrons; electrons move around the nucleus.

**Real-World Reading Link** If you have ever accidentally bitten into a peach pit, you know that your teeth pass easily through the fruit, but cannot dent the hard pit. Similarly, many particles that pass through the outer parts of an atom are deflected by the dense center of the atom.

### The Atom

Many experiments since Dalton's time have proven that atoms do exist. So what exactly is the definition of an atom? To answer this question, consider a gold ring. Suppose you decide to grind the ring down into a pile of gold dust. Each fragment of gold dust still retains all of the properties of gold. If it were possible—which it is not without special equipment—you could continue to divide the gold dust particles into still smaller particles. Eventually, you would encounter a particle that could not be divided any further and still retain the properties of gold. This smallest particle of an element that retains the properties of the element is called an **atom**.

To get an idea of its size, consider the population of the world, which was about  $6.5 \times 10^9$  in 2006. By comparison, a typical solid-copper penny contains  $2.9 \times 10^{22}$  atoms, almost five trillion times the world population! The diameter of a single copper atom is  $1.28 \times 10^{-10}$  m. Placing  $6.5 \times 10^9$  copper atoms side by side would result in a line of copper atoms less than 1 m long. **Figure 4.4** illustrates another way to visualize the size of an atom. Imagine that you increase the size of an atom to be as big as an orange. To keep the proportions between the real sizes of the atom and of the orange, you would have to increase to size of the orange and make it as big as Earth. This illustrates how small atoms are.

■ **Figure 4.4** Imagine that you could increase the size of an atom to make it as big as an orange. At this new scale, an orange would be as big as Earth.




**Connection to Biology** **Looking at atoms** You might think that because atoms are so small, there would be no way to see them. However, an instrument called the scanning tunneling microscope (STM) allows individual atoms to be seen. Just as you need a microscope to study cells in biology, the STM allows you to study atoms. STM work as follows: a fine point is moved above a sample and the interaction of the point with the superficial atoms is recorded electronically. **Figure 4.5** illustrates how individual atoms look when observed with a STM. Scientists are now able to move individual atoms around to form shapes, patterns, and even simple machines. This capability has led to the exciting new field of nanotechnology. The promise of nanotechnology is molecular manufacturing—the atom-by-atom building of machines the size of molecules. As you will read in Chapter 8, a molecule is a group of atoms that are bonded together and act as a unit.

## The Electron


Once scientists were convinced of the existence of atoms, a new set of questions emerged. What is an atom like? Is the composition of an atom uniform throughout, or is it composed of still-smaller particles? Although many scientists researched the atom in the 1800s, it was not until almost 1900 that some of these questions were answered.

**The cathode-ray tube** As scientists tried to unravel the atom, they began to make connections between matter and electric charge. For instance, has your hair ever clung to your comb? To explore the connection, some scientists wondered how electricity might behave in the absence of matter. With the help of the newly invented vacuum pump, they passed electricity through glass tubes from which most of the air had been removed. Such tubes are called cathode-ray tubes.

A typical cathode-ray tube used by researchers for studying the relationship between mass and charge is illustrated in **Figure 4.6**. Note that metal electrodes are located at opposite ends of the tube. The electrode connected to the negative terminal of the battery is called the cathode, and the electrode connected to the positive terminal is called the anode.



■ **Figure 4.5** This image, recorded with a STM, shows the individual atoms of a fatty acid on a graphite surface. The false colors were added later on to improve the contrast between each atom.




■ **Figure 4.6** A cathode-ray tube is a tube with an anode at one end and a cathode at the other end. When a voltage is applied, electricity travels from the cathode to the anode.




■ **Figure 4.7** A tiny hole located in the center of the anode produces a thin beam of electrons. A phosphor coating allows the position of the beam to be determined as it strikes the end of the tube.


**a** Because the cathode ray is deflected in a magnetic field, the particles in the ray must be charged.



**b** Because the cathode ray is deflected toward the positively charged plate by an electric field, the particles in the ray must have a negative charge.



## Real-World Chemistry Cathode Ray



**Television** Television was invented in the 1920's. Conventional television images are formed as cathode rays strike light-producing chemicals that coat the back of the screen.

### FOLDABLES

Incorporate information from this section into your Foldable.

**Sir William Crookes** While working in a darkened laboratory, English physicist Sir William Crookes noticed a flash of light within one of the cathode-ray tubes. A green flash was produced by some form of radiation striking a zinc-sulfide coating that had been applied to the end of the tube. Further work showed that there was a ray (radiation) going through the tube. This ray, originating from the cathode and traveling to the anode, was called a **cathode ray**. The accidental discovery of the cathode ray led to the invention of television. A conventional television is nothing else than a cathode-ray tube.

Scientists continued their research using cathode-ray tubes, and they were fairly convinced by the end of the 1800s of the following:

- Cathode rays were a stream of charged particles.
- The particles carried a negative charge. (The exact value of the negative charge was not known.)

Because changing the metal that makes up the electrodes or varying the gas (at very low pressure) in the cathode-ray tube did not affect the cathode ray produced, researchers concluded that the ray's negative particles were found in all forms of matter. These negatively charged particles that are part of all forms of matter are now known as **electrons**. Some of the experiments used to determine the properties of the cathode ray are shown in **Figure 4.7**.




**Reading Check** Explain how the cathode ray was discovered.

**Mass and charge of the electron** In spite of the progress made from all of the cathode-ray tube experiments, no one succeeded in determining the mass of a single cathode-ray particle. Unable to measure the particle's mass directly, English physicist J. J. Thomson (1856–1940) began a series of cathode-ray tube experiments at Cambridge University in the late 1890s to determine the ratio of its charge to its mass.

**Charge-to-mass ratio** By carefully measuring the effects of both magnetic and electric fields on a cathode ray, Thomson was able to determine the charge-to-mass ratio of the charged particle. He then compared that ratio to other known ratios.

Thomson concluded that the mass of the charged particle was much less than that of a hydrogen atom, the lightest known atom. The conclusion was shocking because it meant there were particles smaller than the atom. In other words, Dalton had been incorrect—atoms were divisible into smaller subatomic particles. Because Dalton's atomic theory had become so widely accepted and Thomson's conclusion was so revolutionary, many other scientists found it hard to accept this new discovery. But Thomson was correct. He had identified the first subatomic particle—the electron. He received a Nobel Prize in 1906 for this discovery.

 **Reading Check** Summarize how Thomson discovered the electron.


### The oil-drop experiment and the charge of an electron

The next significant development came in the early 1910s, when the American physicist Robert Millikan (1868–1953) determined the charge of an electron using the oil-drop apparatus shown in **Figure 4.8**. In this apparatus, oil is sprayed into the chamber above the two parallel charged plates. The top plate has a small hole through which the oil drops. X rays knock out electrons from the air particles between the plates and the electrons stick to the droplets, giving them a negative charge. By varying the intensity of the electric field, Millikan could control the rate of a droplet's fall. He determined that the magnitude of the charge on each drop increased in discrete amounts and determined that the smallest common denominator was  $1.602 \times 10^{-19}$  coulombs. He identified this number as the charge of the electron. This charge was later equated to a single unit of negative charge noted  $1-$ ; in other words, a single electron carries a charge of  $1-$ .

So good was Millikan's experimental setup and technique that the charge he measured almost one hundred years ago is within 1% of the currently accepted value.


**Mass of an electron** Knowing the electron's charge and using the known charge-to-mass ratio, Millikan calculated the mass of an electron. The equation below shows how small the mass of an electron is.

$$\text{Mass of an electron} = 9.1 \times 10^{-28} \text{ g} = \frac{1}{1840} \text{ the mass of a hydrogen atom}$$



**Figure 4.8** The motion of the oil droplets within Millikan's apparatus depends on the charge of droplets and on the electric field. Millikan observed the droplets with the telescope. He could make the droplets fall more slowly, rise, or pause as he varied the strength of the electric field. From his observations, he calculated the charge on each droplet.

■ **Figure 4.9** J. J. Thomson's plum pudding model of the atom states that the atom is a uniform, positively charged sphere containing electrons.




**The plum pudding model** The existence of the electron and the knowledge of some of its properties raised some interesting new questions about the nature of atoms. It was known that matter is neutral—it has no electric charge. You know that matter is neutral from everyday experience: you do not receive an electric shock (except under certain conditions) when you touch an object. If electrons are part of all matter and they possess a negative charge, how can all matter be neutral? Also, if the mass of an electron is so small, what accounts for the rest of the mass in a typical atom?


In an attempt to answer these questions, J. J. Thomson proposed a model of the atom that became known as the plum pudding model. As you can see in **Figure 4.9**, Thomson's model consisted of a spherically shaped atom composed of a uniformly distributed positive charge in which the individual negatively charged electrons resided. As you are about to read, the plum pudding model of the atom did not last for long. **Figure 4.10** summarizes the numerous steps in understanding the structure of the atom.

✓ **Reading Check Explain** why Thomson's model was called the plum pudding model.

### ■ **Figure 4.10** **Development of Modern Atomic Theory**

Current understanding of the properties and behavior of atoms and subatomic particles is based on the work of scientists worldwide during the past two centuries.





■ **Figure 4.11** Based on Thomson's model, Rutherford expected the light alpha particles to pass through gold atoms. He expected only a few of them to be slightly deflected.

## The Nucleus

In 1911, Ernest Rutherford (1871–1937) began to study how positively charged alpha particles (radioactive particles you will read more about later in this chapter) interacted with solid matter. With a small group of scientists, Rutherford conducted an experiment to see if alpha particles would be deflected as they passed through a thin gold foil.

**Rutherford's experiment** In the experiment, a narrow beam of alpha particles was aimed at a thin sheet of gold foil. A zinc-sulfide-coated screen surrounding the gold foil produced a flash of light when struck by an alpha particle. By noting where the flashes occurred, the scientists could determine if the atoms in the gold foil deflected the alpha particles.

Rutherford was aware of Thomson's plum pudding model of the atom. He expected the paths of the massive and fast-moving alpha particles to be only slightly altered by a collision with an electron. And because the positive charge within the gold atoms was thought to be uniformly distributed, he thought it would not alter the paths of the alpha particles, either. **Figure 4.11** shows the results Rutherford expected from the experiment.



1938 Lise Meitner, Otto Hahn, and Fritz Strausman split uranium atoms in a process they called fission.

1954 CERN, the world's largest nuclear physics research center, located in Switzerland, is founded to study particle physics.



2007 The Large Hadron Collider at CERN studies the properties of subatomic particles and nuclear matter.

1939–1945 Scientists in the United States and Germany each work on projects to develop the first atomic weapon.

1968 Scientists provide the first experimental evidence for subatomic particles known as quarks.


Concepts in Motion

**Interactive Time Line** To learn more about these discoveries and others, visit [glencoe.com](http://glencoe.com).






**Interactive Figure** To see an animation of the gold foil experiment, visit [glencoe.com](http://glencoe.com).



■ **Figure 4.12** During Rutherford's experiment, a beam of alpha particles bombarded a thin gold foil. Most of the alpha particles went through the gold foil. However, a few of them bounced back, some at large angles.

■ **Figure 4.13** In Rutherford's nuclear model, the atom is composed of a dense, positively charged nucleus that is surrounded by negative electrons. Alpha particles passing far from the nucleus are only slightly deflected. Alpha particles directly approaching the nucleus are deflected at large angles.

**Infer** what force causes the deflection of alpha particles.



The actual results observed by Rutherford and his colleagues are shown in **Figure 4.12**. A few of the alpha particles were deflected at large angles. Several particles were deflected straight back toward the source. Rutherford likened the results to firing a large artillery shell at a sheet of paper and the shell coming back at the cannon.

**Rutherford's model of the atom** Rutherford concluded that the plum pudding model was incorrect because it could not explain the results of the gold foil experiment. Considering the properties of the alpha particles and the electrons, and the frequency of the deflections, he calculated that an atom consisted mostly of empty space through which the electrons move. He also concluded that almost all of the atom's positive charge and almost all of its mass were contained in a tiny, dense region in the center of the atom, which he called the **nucleus**. The negatively charged electrons are held within the atom by their attraction to the positively charged nucleus. Rutherford's nuclear atomic model is shown in **Figure 4.13**.

Because the nucleus occupies such a small space and contains most of an atom's mass, it is incredibly dense. If a nucleus were the size of the dot in the exclamation point at the end of this sentence, its mass would be approximately as much as that of 70 automobiles! The volume of space through which the electrons move is huge compared to the volume of the nucleus. A typical atom's diameter is approximately 10,000 times the diameter of the nucleus. If an atom had a diameter of two football fields, the nucleus would be the size of a nickel.

**Reading Check** Describe Rutherford's model of the atom.

The repulsive force produced between the positive nucleus and the positive alpha particles causes the deflections. **Figure 4.13** illustrates how Rutherford's nuclear atomic model explained the results of the gold foil experiment. The nuclear model also explains the neutral nature of matter: the positive charge of the nucleus balances the negative charge of the electrons. However, the model still could not account for all of the atom's mass.

**The proton and the neutron** By 1920, Rutherford had refined the concept of the nucleus and concluded that the nucleus contained positively charged particles called protons. A **proton** is a subatomic particle carrying a charge equal to but opposite that of an electron; that is, a proton has a charge of  $1+$ . In 1932, Rutherford's coworker, English physicist James Chadwick (1891–1974), showed that the nucleus also contained another subatomic neutral particle, called the neutron. A **neutron** is a subatomic particle that has a mass nearly equal to that of a proton, but it carries no electric charge. In 1935, Chadwick received the Nobel Prize in Physics for proving the existence of neutrons.

## VOCABULARY

### SCIENCE USAGE V. COMMON USAGE

#### Neutral

**Science usage:** to have no electric charge

*Neutrons have a charge of zero. They are neutral particles.*

**Common usage:** not engaged in either side

*Switzerland remained neutral during World War II.*

## DATA ANALYSIS LAB

Based on Real Data\*

### Interpret Scientific Illustrations



**What are the apparent atomic distances of carbon atoms in a well-defined crystalline material?** To visualize individual atoms, a group of scientists used a scanning tunneling microscope (STM) to test a crystalline material called highly ordered pyrolytic graphite (HOPG). An STM is an instrument used to perform surface atomic-scale imaging.

#### Data and Observations

The image shows all of the carbon atoms in the surface layer of the graphite material. Each hexagonal ring, indicated by the drawing in the figure, consists of three brighter spots separated by three fainter spots. These bright spots are from alternate carbon atoms in the surface layer of the graphite structure. The cross-sectional view below the photo corresponds to the line drawn in the image. It indicates the atomic periodicity and apparent atomic distances.


#### Think Critically

1. **Estimate** the distance between two nearest bright spots.
2. **Estimate** the distance between two nearest neighbor spots (brighter–fainter, marked with triangles in the figure).



\*Data obtained from: Chaun-Jian Zhong et al. 2003. Atomic scale imaging: a hands-on scanning probe microscopy laboratory for undergraduates. *Journal of Chemical Education* 80: 194–197.

3. **State** What do the black spots in the image represent?
4. **Explain** How many carbon atoms are across the line drawn in the image?



■ **Figure 4.14** Atoms are composed of a nucleus containing protons and neutrons, and surrounded by a cloud of electrons.

## Concepts In Motion

**Interactive Figure** To see an animation of the structure of the atom, visit [glencoe.com](http://glencoe.com).

**Table 4.3** Properties of Subatomic Particles

**Interactive Table** Explore the properties of subatomic particles at [glencoe.com](http://glencoe.com).

Particle	Symbol	Location	Relative Electric Charge	Relative Mass	Actual Mass (g)
Electron	$e^-$	In the space surrounding the nucleus	1-	$\frac{1}{1840}$	$9.11 \times 10^{-28}$
Proton	p	In the nucleus	1+	1	$1.673 \times 10^{-24}$
Neutron	n	In the nucleus	0	1	$1.675 \times 10^{-24}$

**Completing the model of the atom** All atoms are made up of the three fundamental subatomic particles—the electron, the proton, and the neutron. Atoms are spherically shaped, with a small, dense nucleus of positive charge surrounded by one or more negatively charged electrons. Most of an atom consists of fast-moving electrons traveling through the empty space surrounding the nucleus. The electrons are held within the atom by their attraction to the positively charged nucleus. The nucleus, which is composed of neutral neutrons (hydrogen’s single-proton nucleus is an exception) and positively charged protons, contains all of an atom’s positive charge and more than 99.97% of its mass. It occupies only about one ten-thousandth of the volume of the atom. Because an atom is electrically neutral, the number of protons in the nucleus equals the number of electrons surrounding the nucleus. The features of a typical atom are shown in **Figure 4.14**, and the properties of the fundamental subatomic particles are summarized in **Table 4.3**.

Subatomic particle research is still a major interest to modern scientists. In fact, scientists have determined that protons and neutrons have their own structures. They are composed of subatomic particles called quarks. These particles will not be covered in this textbook because scientists do not yet understand if or how they affect chemical behavior. As you will learn in later chapters, chemical behavior can be explained by considering only an atom’s electrons.

## Section 4.2 Assessment

### Section Summary

- ▶ An atom is the smallest particle of an element that maintains the properties of that element.
- ▶ Electrons have a 1– charge, protons have a 1+ charge, and neutrons have no charge.
- ▶ An atom consists mostly of empty space surrounding the nucleus.

7. **MAIN Idea** Describe the structure of a typical atom. Identify where each subatomic particle is located.
8. **Compare and contrast** Thomson’s plum pudding atomic model with Rutherford’s nuclear atomic model.
9. **Evaluate** the experiments that led to the conclusion that electrons are negatively charged particles found in all matter.
10. **Compare** the relative charge and mass of each of the subatomic particles.
11. **Calculate** What is the difference expressed in kilograms between the mass of a proton and the mass of an electron?

## Section 4.3

### Objectives

- **Explain** the role of atomic number in determining the identity of an atom.
- **Define** an isotope.
- **Explain** why atomic masses are not whole numbers.
- **Calculate** the number of electrons, protons, and neutrons in an atom given its mass number and atomic number.

### Review Vocabulary

**periodic table:** a chart that organizes all known elements into a grid of horizontal rows (periods) and vertical columns (groups or families) arranged by increasing atomic number

### New Vocabulary

atomic number  
isotope  
mass number  
atomic mass unit (amu)  
atomic mass

## How Atoms Differ

**MAIN Idea** The number of protons and the mass number define the type of atom.

**Real-World Reading Link** You are probably aware that numbers are used every day to identify people and objects. For example, people can be identified by their Social Security numbers and computers by their IP addresses. Atoms and nuclei are also identified by numbers.

### Atomic Number

As shown in the periodic table of the elements inside the back cover of this textbook, there are more than 110 different elements. What makes an atom of one element different from an atom of another element?

Not long after Rutherford's gold foil experiment, the English scientist Henry Moseley (1887–1915) discovered that atoms of each element contain a unique positive charge in their nuclei. Thus, the number of protons in an atom identifies it as an atom of a particular element. The number of protons in an atom is referred to as the **atomic number**. The information provided by the periodic table for hydrogen is shown in **Figure 4.15**. The number 1 above the symbol for hydrogen (H) is the number of protons, or the atomic number. Moving across the periodic table to the right, you will next come to helium (He). It has two protons in its nucleus, and thus it has an atomic number of 2. The next row begins with lithium (Li), atomic number 3, followed by beryllium (Be), atomic number 4, and so on. The periodic table is organized left-to-right and top-to-bottom by increasing atomic number.

Because all atoms are neutral, the number of protons and electrons in an atom must be equal. Thus, once you know the atomic number of an element, you know the number of protons and the number of electrons an atom of that element contains. For example, an atom of lithium, atomic number 3, contains three protons and three electrons.

### Atomic number

$$\text{atomic number} = \text{number of protons} \\ = \text{number of electrons}$$

The atomic number of an atom equals its number of protons and its number of electrons.

- **Figure 4.15** In the periodic table, each element is represented by its chemical name, atomic number, chemical symbol, and average atomic mass.  
**Determine** the number of protons and the number of electrons in an atom of gold.

Hydrogen	Chemical name
1	Atomic number
H	Chemical symbol
1.008	Average atomic mass



## EXAMPLE Problem 4.1

### Math Handbook

Solving Algebraic Equations  
pages 954–955

**Atomic Number** Complete the following table.

Composition of Several Elements				
	Element	Atomic Number	Protons	Electrons
a.	Pb	82		
b.			8	
c.				30

### 1 Analyze the Problem

Apply the relationship among atomic number, number of protons, and number of electrons to complete most of the table. Then, use the periodic table to identify the element.

#### Known

- a. element = Pb, atomic number = 82
- b. number of protons = 8
- c. number of electrons = 30

#### Unknown

- a. number of protons ( $N_p$ ), number of electrons ( $N_e$ ) = ?
- b. **element**, atomic number ( $Z$ ),  $N_e$  = ?
- c. **element**,  $Z$ ,  $N_p$  = ?

### 2 Solve for the Unknown

- a. number of protons = atomic number

$$N_p = 82$$

number of electrons = number of protons

$$N_e = 82$$

The number of protons and the number of electrons is 82.

- b. atomic number = number of protons

$$Z = 8$$

number of electrons = number of protons

$$N_e = 8$$

The atomic number and the number of electrons is 8.

The **element** is **oxygen (O)**.

- c. number of protons = number of electrons

$$N_p = 30$$

atomic number = number of protons

$$Z = 30$$

The atomic number and the number of protons is 30.

The **element** is **zinc (Zn)**.

Apply the atomic-number relationship.

Substitute atomic number = 82.

Apply the atomic-number relationship.

Substitute number of protons = 8.

Consult the periodic table to identify the element.

Apply the atomic-number relationship.

Substitute number of electrons = 30.

Consult the periodic table to identify the element.

### 3 Evaluate the Answer

The answers agree with atomic numbers and element symbols given in the periodic table.

## PRACTICE Problems

Extra Practice Pages 977–978 and [glencoe.com](http://glencoe.com)

- 12. How many protons and electrons are in each atom?
  - a. radon
  - b. magnesium
- 13. An atom of an element contains 66 electrons. Which element is it?
- 14. An atom of an element contains 14 protons. Which element is it?
- 15. **Challenge** Do the atoms shown in the figure to the right have the same atomic number?

$9e^-$

$10n$

$9p$   
 $9n$

## Isotopes and Mass Number

Dalton was incorrect about atoms being indivisible and in stating that all atoms of an element are identical. All atoms of an element have the same number of protons and electrons, but the number of neutrons might differ. For example, there are three types of potassium atoms that occur naturally. All three types contain 19 protons and 19 electrons. However, one type of potassium atom contains 20 neutrons, another 21 neutrons, and still another 22 neutrons. Atoms with the same number of protons but different numbers of neutrons are called **isotopes**.

**Mass of isotopes** Isotopes containing more neutrons have a greater mass. In spite of these differences, isotopes of an atom have the same chemical behavior. As you will read later in this textbook, chemical behavior is determined only by the number of electrons an atom has.

**Isotope notation** Each isotope of an element is identified with a number called the mass number. The **mass number** is the sum of the atomic number (or number of protons) and neutrons in the nucleus.


### Mass number

$$\text{mass number} = \text{atomic number} + \text{number of neutrons}$$

The mass number of an atom is the sum of its atomic number and its number of neutrons.


For example, copper has two isotopes. The isotope with 29 protons and 34 neutrons has a mass number of 63 ( $29 + 34 = 63$ ), and is called copper-63 (also written  $^{63}\text{Cu}$  or Cu-63). The isotope with 29 protons and 35 neutrons is called copper-65. Chemists often write out isotopes using a notation involving the chemical symbol, atomic number, and mass number, as shown in **Figure 4.16**.

**Natural abundance of isotopes** In nature, most elements are found as mixtures of isotopes. Usually, no matter where a sample of an element is obtained, the relative abundance of each isotope is constant. For example, in a banana, 93.26% of the potassium atoms have 20 neutrons, 6.73% have 22 neutrons, and 0.01% have 21 neutrons. In another banana, or in a different source of potassium, the percentage composition of the potassium isotopes will still be the same. The three potassium isotopes are summarized in **Figure 4.17**.



■ **Figure 4.16** Cu is the chemical symbol for copper. Copper, which was used to make this Chinese gong, is composed of 69.2% copper-63 and 30.8% copper-65.

	Potassium-39	Potassium-40	Potassium-41
Protons	19	19	19
Neutrons	20	21	22
Electrons	19	19	19



■ **Figure 4.17** Potassium has three naturally occurring isotopes: potassium-39, potassium-40, and potassium-41.

**List the number of protons, neutrons, and electrons in each potassium isotope.**

## EXAMPLE Problem 4.2

**Use Atomic Number and Mass Number** A chemistry laboratory has analyzed the composition of isotopes of several elements. The composition data is given in the table below. Determine the number of protons, electrons, and neutrons in the isotope of neon. Name the isotope and give its symbol.

Isotope Composition Data			
	Element	Atomic Number	Mass Number
a.	Neon	10	22
b.	Calcium	20	46
c.	Oxygen	8	17
d.	Iron	26	57
e.	Zinc	30	64
f.	Mercury	80	204

### 1 Analyze the Problem

You are given some data for neon in the table. The symbol for neon can be found on the periodic table. From the atomic number, the number of protons and electrons in the isotope are known. The number of neutrons in the isotope can be found by subtracting the atomic number from the mass number.

#### Known

element: neon  
atomic number = 10  
mass number = 22

#### Unknown

number of protons ( $N_p$ ), electrons ( $N_e$ ), and neutrons ( $N_n$ ) = ?  
name of isotope = ?  
symbol for isotope = ?

### 2 Solve for the Unknown

number of protons = atomic number = **10**  
number of electrons = atomic number = **10**  
number of neutrons = mass number – atomic number

$$N_n = 22 - 10 = \mathbf{12}$$

The **name** of the isotope is **neon-22**.

The **symbol** for the isotope is  ${}^{22}_{10}\text{Ne}$ .

Apply the atomic number relationship.

Use the atomic number and the mass number to calculate the number of neutrons.

Substitute mass number = 22 and atomic number = 10

Use the element name and mass number to write the isotope's name.

Use the chemical symbol, mass number, and atomic number to write out the isotope in symbolic notation form.

### 3 Evaluate the Answer

The relationships among number of electrons, protons, and neutrons have been applied correctly. The isotope's name and symbol are in the correct format. Refer to pages 944–945 the Elements Handbook to learn more about neon.

## PRACTICE Problems

Extra Practice Page 978 and [glencoe.com](http://glencoe.com)

- Determine the number of protons, electrons, and neutrons for isotopes **b.–f.** in the table above. Name each isotope, and write its symbol.
- Challenge** An atom has a mass number of 55. Its number of neutrons is the sum of its atomic number and five. How many protons, neutrons, and electrons does this atom have? What is the identity of this atom?

Table 4.4		Masses of Subatomic Particles
Particle	Mass (amu)	
Electron	0.000549	
Proton	1.007276	
Neutron	1.008665	

## Mass of Atoms

Recall from **Table 4.3** that the masses of both protons and neutrons are approximately  $1.67 \times 10^{-24}$  g. While this is a small mass, the mass of an electron is even smaller—only about 1/1840 that of a proton or a neutron.

**Atomic mass unit** Because these extremely small masses expressed in scientific notation are difficult to work with, chemists have developed a method of measuring the mass of an atom relative to the mass of a specific atomic standard. That standard is the carbon-12 atom. Scientists assigned the carbon-12 atom a mass of exactly 12 atomic mass units. Thus, one **atomic mass unit (amu)** is defined as one-twelfth the mass of a carbon-12 atom. Although a mass of 1 amu is nearly equal to the mass of a single proton or a single neutron, it is important to realize that the values are slightly different. **Table 4.4** gives the masses of the subatomic particles in terms of amu.

**Atomic mass** Because an atom's mass depends mainly on the number of protons and neutrons it contains, and because protons and neutrons have masses close to 1 amu, you might expect the atomic mass of an element to always be nearly a whole number. However, this is often not the case. The explanation involves how atomic mass is defined. The **atomic mass** of an element is the weighted average mass of the isotopes of that element. Because isotopes have different mass, the weighted average is not a whole number. The calculation of the atomic mass of chlorine is illustrated in **Figure 4.18**.

## VOCABULARY

### ACADEMIC VOCABULARY

#### Specific

characterized by precise formulation or accurate restriction


*Some diseases have specific symptoms.*

■ **Figure 4.18** To calculate the weighted average atomic mass of chlorine, you first need to calculate the mass contribution of each isotope.



**Personal Tutor** For an online tutorial on finding an average, visit [glencoe.com](http://glencoe.com).

### Calculate the Weighted Average Atomic Mass of Chlorine







■ **Figure 4.19** Bromine is extracted from sea water and salt lakes. The Dead Sea area in Israel is one of the major bromine production sites in the world. Applications of bromine include microbe and algae control in swimming pools and flame-retardants. It is also used in medicines, oils, paints, and pesticides.

Chlorine exists naturally as a mixture of about 76% chlorine-35 and 24% chlorine-37. It has an atomic mass of 35.453 amu. Because atomic mass is a weighted average, the chlorine-35 atoms, which exist in greater abundance than the chlorine-37 atoms, have a greater effect in determining the atomic mass. The atomic mass of chlorine is calculated by multiplying each isotope's percent abundance by its atomic mass and then adding the products. The process is similar to calculating an average grade. You can calculate the atomic mass of any element if you know the number of naturally occurring isotopes, their masses, and their percent abundances.



**Reading Check** Explain how to calculate atomic mass.

**Isotope abundances** Analyzing an element's mass can indicate the most abundant isotope for that element. For example, fluorine (F) has an atomic mass that is extremely close to 19 amu. If fluorine had several fairly abundant isotopes, its atomic mass would not likely be so close to a whole number. Thus, you might conclude that all naturally occurring fluorine is probably in the form of fluorine-19 ( $^{19}\text{F}$ ). Indeed, 100% of naturally occurring fluorine is in the form of fluorine-19. While this type of reasoning generally works well, it is not foolproof. Consider bromine (Br). It has an atomic mass of 79.904 amu. With a mass so close to 80 amu, it seems likely that the most common bromine isotope would be bromine-80. However, Bromine's two isotopes are bromine-79 (78.918 amu, 50.69%) and bromine-81 (80.917 amu, 49.31%). There is no bromine-80 isotope. **Figure 4.19** shows one of the major production sites of bromine, located in the Dead Sea area. Refer to page 940 of the Elements Handbook to learn more about chlorine, fluorine, and bromine.

## MiniLab

### Model Isotopes

**How can you calculate the atomic mass of an element using the percentage abundance of its isotopes?** Because they have different compositions, pre- and post-1982 pennies can be used to model an element with two naturally occurring isotopes. From the penny 'isotope' data, you can determine the mass of each penny isotope and the average mass of a penny.

#### Procedure

1. Read and complete the lab safety form.
2. Get a bag of **pennies** from your teacher, and sort the pennies by date into two groups: pre-1982 pennies and post-1982 pennies. Count and record the total number of pennies and the number in each group.
3. Using a **balance**, determine the mass of 10 pennies from each group. Record each mass to the nearest 0.01 g. Divide the total mass of each group by 10 to get the average mass of a pre- and post-1982 penny isotope.

#### Analysis

1. **Calculate** the percentage abundance of each group using data from Step 2. To do this, divide the number of pennies in each group by the total number of pennies.
2. **Determine** the atomic mass of a penny using the percentage abundance of each "isotope" and data from Step 3. To do this, use the following equation:  
$$\text{mass contribution} = (\% \text{ abundance})(\text{mass})$$
Total the mass contributions to determine the atomic mass. Remember that the percent abundance is a percentage.
3. **Infer** whether the atomic mass would be different if you received another bag of pennies containing a different mixture of pre- and post-1982 pennies. Explain your reasoning.
4. **Explain** why the average mass of each type of penny was determined by measuring 10 pennies instead of by measuring and using the mass of a single penny from each group.

## EXAMPLE Problem 4.3

**Calculate Atomic Mass** Given the data in the table, calculate the atomic mass of unknown Element X. Then, identify the unknown element, which is used medically to treat some mental disorders.

### 1 Analyze the Problem

Calculate the atomic mass and use the periodic table to confirm.

#### Known

${}^6\text{X}$ : mass = 6.015 amu  
abundance = 7.59% = 0.0759  
 ${}^7\text{X}$ : mass = 7.016 amu  
abundance = 92.41% = 0.9241

#### Unknown

atomic mass of X = ? amu

element X = ?

Isotope Abundance for Element X

Isotope	Mass (amu)	Percent Abundance
${}^6\text{X}$	6.015	7.59%
${}^7\text{X}$	7.016	92.41%

### 2 Solve for the Unknown

${}^6\text{X}$ : mass contribution = (mass)(percent abundance)  
mass contribution = (6.015 amu)(0.0759) = 0.4565 amu  
 ${}^7\text{X}$ : mass contribution = (mass)(percent abundance)  
mass contribution = (7.016 amu)(0.9241) = 6.483 amu

atomic mass of X = (0.4565 amu + 6.483 amu) = **6.939 amu**

The **element** with a mass 6.939 amu is **lithium (Li)**.

Calculate  ${}^6\text{X}$ 's contribution.

Substitute mass = 6.015 amu and abundance = 0.0759.

Calculate  ${}^7\text{X}$ 's contribution.

Substitute mass = 7.016 amu and abundance = 0.9241.

Total the mass contributions to find the atomic mass.

Identify the element using the periodic table.

### 3 Evaluate the Answer

The result of the calculation agrees with the atomic mass given in the periodic table. The masses of the isotopes have four significant figures, so the atomic mass is also expressed with four significant figures. Refer to the Elements Handbook to learn more about lithium.

## PRACTICE Problems

Extra Practice Page 978 and [glencoe.com](http://glencoe.com)

18. Boron (B) has two naturally occurring isotopes: boron-10 (abundance = 19.8%, mass = 10.013 amu) and boron-11 (abundance = 80.2%, mass = 11.009 amu). Calculate the atomic mass of boron.
19. **Challenge** Nitrogen has two naturally occurring isotopes, N-14 and N-15. Its atomic mass is 14.007. Which isotope is more abundant? Explain your answer.

## Section 4.3 Assessment

### Section Summary

- The atomic number of an atom is given by its number of protons. The mass number of an atom is the sum of its neutrons and protons.
- Atoms of the same element with different numbers of neutrons are called isotopes.
- The atomic mass of an element is a weighted average of the masses of all of its naturally occurring isotopes.
- **20. MAIN Idea** Explain how the type of an atom is defined.
- **21. Recall** Which subatomic particle identifies an atom as that of a particular element?
- **22. Explain** how the existence of isotopes is related to the fact that atomic masses are not whole numbers.
- **23. Calculate** Copper has two isotopes: Cu-63 (abundance = 69.2%, mass = 62.930 amu) and Cu-65 (abundance = 30.8%, mass = 64.928 amu). Calculate the atomic mass of copper.
- **24. Calculate** Three magnesium isotopes have atomic masses and relative abundances of 23.985 amu (79.99%), 24.986 amu (10.00%), and 25.982 (11.01%). Calculate the atomic mass of magnesium.

## Section 4.4

### Objectives

- **Explain** the relationship between unstable nuclei and radioactive decay.
- **Characterize** alpha, beta, and gamma radiation in terms of mass and charge.

### Review Vocabulary

**element:** a pure substance that cannot be broken down into simpler substances by physical or chemical means

### New Vocabulary

radioactivity  
radiation  
nuclear reaction  
radioactive decay  
alpha radiation  
alpha particle  
nuclear equation  
beta radiation  
beta particle  
gamma ray

# Unstable Nuclei and Radioactive Decay

**MAIN Idea** Unstable atoms emit radiation to gain stability.

**Real-World Reading Link** Try dropping a rock from the height of your waist. The rock goes from a higher energy state at your waist to a lower energy state on the floor. A similar process happens with nuclei in an unstable state.

## Radioactivity

Recall from Chapter 3 that a chemical reaction is the change of one or more substances into new substances and involves only an atom's electrons. Although atoms might be rearranged, their identity remains the same. Another type of reaction, called a nuclear reaction, can change an element into a new element.

**Nuclear reactions** In the late 1890s, scientists noticed that some substances spontaneously emitted radiation in a process they named **radioactivity**. The rays and particles emitted by the radioactive material were called **radiation**. Scientists discovered that radioactive atoms undergo changes that can alter their identities. A reaction that involves a change in an atom's nucleus is called a **nuclear reaction**. The discovery of these nuclear reactions was a major breakthrough, as no chemical reaction had ever resulted in the formation of new kinds of atoms.

Radioactive atoms emit radiation because their nuclei are unstable. Unstable systems, whether they are atoms or people doing handstands, as shown in **Figure 4.20**, gain stability by losing energy.

**Radioactive decay** Unstable nuclei lose energy by emitting radiation in a spontaneous process called **radioactive decay**. Unstable atoms undergo radioactive decay until they form stable atoms, often of a different element. Just as a rock loses gravitational potential energy and reaches a stable state when falling to the ground, an atom can lose energy and reach a stable state when emitting radiation.


■ **Figure 4.20** Being in a handstand position is an unstable state. Like unstable atoms, people doing handstands eventually return to a more stable state—standing on their feet—by losing potential energy.



## Types of Radiation


Scientists began researching radioactivity in the late 1800s. They investigated the effect of electric fields on radiation. By directing radiation from a radioactive source between two electrically charged plates, scientists were able to identify three different types of radiation based on their electric charge. As shown in **Figure 4.21**, radiation were deflected toward the negative plate, the positive plate, or not at all.

**Alpha radiation** The radiation that was deflected toward the negatively charged plate was named **alpha radiation**. It is made up of alpha particles. An **alpha particle** contains two protons and two neutrons, and thus has a 2+ charge, which explains why alpha particles are attracted to the negatively charged plate as shown in **Figure 4.21**. An alpha particle is equivalent to a helium-4 nucleus and is represented by  ${}^4_2\text{He}$  or  $\alpha$ . The alpha decay of radioactive radium-226 into radon-222 is shown below.



Note that a new element, radon (Rn), is created as a result of the alpha decay of the unstable radium-226 nucleus. The type of equation shown above is known as a **nuclear equation**. It shows the atomic numbers and mass numbers of the particles involved. The mass number is conserved in nuclear equations.

**Beta radiation** The radiation that was deflected toward the positively charged plate was named **beta radiation**. This radiation consists of fast-moving beta particles. Each **beta particle** is an electron with a 1- charge. The negative charge of the beta particle explains why it is attracted to the positively charged plate shown in **Figure 4.21**. Beta particles are represented by the symbol  $\beta$  or  $e^-$ . The beta decay of carbon-14 into nitrogen-14 is shown below. The beta decay of unstable carbon-14 results in the creation of the new atom, nitrogen (N).




### CAREERS IN CHEMISTRY

**Chemistry Teacher** Chemistry teachers work in high schools and colleges. They lecture, guide discussions, conduct experiments, supervise lab work, and lead field trips. High school teachers might also be asked to monitor study halls and serve on committees. College instructors might be required to do research and publish their findings. For more information on chemistry careers, visit [glencoe.com](http://glencoe.com).

■ **Figure 4.21** An electric field will deflect radiation in different directions, depending on the electric charge of the radiation.

**Explain** why beta particles are deflected toward the positive plate, alpha particles are deflected toward the negative plate, and gamma rays are not deflected.




**Interactive Figure** To see an animation of radiation deflection, visit [glencoe.com](http://glencoe.com).



Table 4.5		Characteristics of Radiation		
	Alpha	Beta	Gamma	
Symbol	${}^4_2\text{He}$ or $\alpha$	$e^-$ or $\beta$	$\gamma$	
Mass (amu)	4	$\frac{1}{1840}$	0	
Mass (kg)	$6.65 \times 10^{-27}$	$9.11 \times 10^{-31}$	0	
Charge	2+	1-	0	

**Gamma radiation** The third common type of radiation is called gamma radiation, or gamma rays. A **gamma ray** is a high-energy radiation that possesses no mass and is denoted by the symbol  $\gamma$ . Because they are neutral, gamma rays are not deflected by electric or magnetic fields. They usually accompany alpha and beta radiation, and they account for most of the energy lost during radioactive decays. For example, gamma rays accompany the alpha decay of uranium-238.



Because gamma rays are massless, the emission of gamma rays by themselves cannot result in the formation of a new atom. **Table 4.5** summarizes the basic characteristics of alpha, beta, and gamma radiation.

**Nuclear stability** The primary factor in determining an atom's stability is its ratio of neutrons to protons. Atoms that contain either too many or too few neutrons are unstable and lose energy through radioactive decay to form a stable nucleus. They emit alpha and beta particles and these emissions affect the neutron-to-proton ratio of the newly created nucleus. Eventually, radioactive atoms undergo enough radioactive decay to form stable, nonradioactive atoms. This topic will be covered in detail in Chapter 24.

## Section 4.4 Assessment

### Section Summary

- ▶ Chemical reactions involve changes in the electrons surrounding an atom. Nuclear reactions involve changes in the nucleus of an atom.
- ▶ There are three types of radiation: alpha (charge of 2+), beta (charge of 1-), and gamma (no charge).
- ▶ The neutron-to-proton ratio of an atom's nucleus determines its stability.


- 25. **MAIN Idea** Explain how unstable atoms gain stability.
- 26. **State** what quantities are conserved when balancing a nuclear reaction.
- 27. **Classify** each of the following as a chemical reaction, a nuclear reaction, or neither.
  - a. Thorium emits a beta particle.
  - b. Two atoms share electrons to form a bond.
  - c. A sample of pure sulfur emits heat energy as it slowly cools.
  - d. A piece of iron rusts.
- 28. **Calculate** How much heavier is an alpha particle than an electron?
- 29. **Create** a table showing how each type of radiation affects the atomic number and the mass number of an atom.

# HOW IT WORKS



## Mass Spectrometer: Chemical Detective

Imagine a forensic scientist needs to identify the inks used on a document to test for possible counterfeiting. The scientist can analyze the inks using a mass spectrometer, such as the one shown at left. A mass spectrometer breaks the compounds in a sample of an unknown substance into smaller fragments. The fragments are then separated according to their masses, and the exact composition of the sample can be determined. Mass spectrometry is one of the most important techniques for studying unknown substances.



### WRITING in Chemistry

**Summarize** Research a case in which a mass spectrometer was used to distinguish between different types of ink, and write a summary of the procedure and results. Visit [glencoe.com](http://glencoe.com) to learn more about mass spectrometers.

# CHEMLAB

## MODEL ATOMIC MASS

**Background:** Most elements in nature occur as a mixture of isotopes. The weighted average atomic mass of an element can be determined from the atomic mass and the relative abundance of each isotope. In this activity, you will model the isotopes of the imaginary element “Snackium.” The measurements you make will be used to calculate a weighted average mass that represents the average atomic mass of “Snackium.”

**Question:** How are the atomic masses of the natural isotopic mixtures calculated?

### Materials

balance  
calculator  
bag of snack mix

### Safety Precautions



**WARNING:** Do not eat food used in lab work.

### Procedure

1. Read and complete the lab safety form.
2. Create a table to record your data. The table will contain the mass and the abundance of each type of snack present in the mixture.
3. Open your snack-mix bag. Handle the pieces with care.
4. Organize the snack pieces into groups based on their types.
5. Count the number of snack pieces in each of your groups.
6. Record the number of snack pieces in each group and the total number of snack pieces in your data table.
7. Measure the mass of one piece from each group and record the mass in your data table.
8. **Cleanup and Disposal** Dispose of the snack pieces as directed by your teacher. Return all lab equipment to its designated location.

### Analyze and Conclude

1. **Calculate** Find the percent abundance of the pieces by dividing the individual-piece quantity by the total number of snack pieces.



2. **Calculate** Use the isotopic percent abundance of the snack pieces and the mass to calculate the weighted average atomic mass for your element “Snackium.”
3. **Interpret** Explain why the weighted average atomic mass of the element “Snackium” is not equal to the mass of any of the pieces.
4. **Peer Review** Gather the average atomic mass data from other lab groups. Explain any differences between your data and the data obtained by other groups.
5. **Apply** Why are the atomic masses on the periodic table not expressed as whole numbers like the mass number of an element?
6. **Research** Look in a chemical reference book to determine whether all elements in the periodic table have isotopes. What is the range of the number of isotopes chemical elements have?
7. **Error Analysis** What sources of error could have led the lab groups to different final values? What modifications could you make in this investigation to reduce the incidence of error?

### INQUIRY EXTENSION

**Predict** Based on your experience in this lab, look up the atomic masses of several elements on the periodic table and predict the most abundant isotope for each element.

**BIG Idea** Atoms are the fundamental building blocks of matter.

### Section 4.1 Early Ideas About Matter

**MAIN Idea** The ancient Greeks tried to explain matter, but the scientific study of the atom began with John Dalton in the early 1800s.**Vocabulary**

- Dalton's atomic theory (p. 104)

**Key Concepts**

- Democritus was the first person to propose the existence of atoms.
- According to Democritus, atoms are solid, homogeneous, and indivisible.
- Aristotle did not believe in the existence of atoms.
- John Dalton's atomic theory is based on numerous scientific experiments.

### Section 4.2 Defining the Atom

**MAIN Idea** An atom is made of a nucleus containing protons and neutrons; electrons move around the nucleus.**Vocabulary**

- atom (p. 106)
- cathode ray (p. 108)
- electron (p. 108)
- neutron (p. 113)
- nucleus (p. 112)
- proton (p. 113)

**Key Concepts**

- An atom is the smallest particle of an element that maintains the properties of that element.
- Electrons have a 1- charge, protons have a 1+ charge, and neutrons have no charge.
- An atom consists mostly of empty space surrounding the nucleus.

### Section 4.3 How Atoms Differ

**MAIN Idea** The number of protons and the mass number define the type of atom.**Vocabulary**

- atomic mass (p. 119)
- atomic mass unit (amu) (p. 119)
- atomic number (p. 115)
- isotope (p. 117)
- mass number (p. 117)

**Key Concepts**

- The atomic number of an atom is given by its number of protons. The mass number of an atom is the sum of its neutrons and protons.

$$\text{atomic number} = \text{number of protons} = \text{number of electrons}$$

$$\text{mass number} = \text{atomic number} + \text{number of neutrons}$$

- Atoms of the same element with different numbers of neutrons are called isotopes.
- The atomic mass of an element is a weighted average of the masses of all of its naturally occurring isotopes.

### Section 4.4 Unstable Nuclei and Radioactive Decay

**MAIN Idea** Unstable atoms emit radiation to gain stability.**Vocabulary**

- alpha particle (p. 123)
- alpha radiation (p. 123)
- beta particle (p. 123)
- beta radiation (p. 123)
- gamma ray (p. 124)
- nuclear equation (p. 123)
- nuclear reaction (p. 122)
- radioactivity (p. 122)
- radiation (p. 122)
- radioactive decay (p. 122)

**Key Concepts**

- Chemical reactions involve changes in the electrons surrounding an atom. Nuclear reactions involve changes in the nucleus of an atom.
- There are three types of radiation: alpha (charge of 2+), beta (charge of 1-), and gamma (no charge).
- The neutron-to-proton ratio of an atom's nucleus determines its stability.



## Section 4.1


## Mastering Concepts

30. Who originally proposed the concept that matter is composed of tiny, indivisible particles?
31. Whose work is credited with being the beginning of modern atomic theory?
32. Distinguish between Democritus's ideas and Dalton's atomic theory.
33. **Ideas and Scientific Methods** Was Democritus's proposal of the existence of atoms based on scientific methods or ideas? Explain.
34. Explain why Democritus was unable to experimentally verify his ideas.
35. What was Aristotle's objection to the atomic theory?
36. State the main points of Dalton's atomic theory using your own words. Which parts of Dalton's theory were later found to be erroneous? Explain why.
37. **Conservation of Mass** Explain how Dalton's atomic theory offered a convincing explanation of the observation that mass is conserved in chemical reactions.
38. Define *matter* and give two everyday examples.

## Section 4.2

## Mastering Concepts


39. What particles are found in the nucleus of an atom? What is the charge of the nucleus?
40. How was the overall charge distributed in the plum pudding model?
41. How did the charge distribution in the plum pudding model affect alpha particles passing through an atom?



■ Figure 4.22

42. Label the subatomic particles shown in Figure 4.22.
43. Arrange the following subatomic particles in order of increasing mass: neutron, electron, and proton.

44. Explain why atoms are electrically neutral.
45. What is the charge of the nucleus of element 89?
46. Which particles account for most of an atom's mass?
47. If you had a balance that could determine the mass of a proton, how many electrons would you need to weigh on the same balance to measure the same mass as that of a single proton?
48. **Cathode-Ray Tubes** Which subatomic particle was discovered by researchers working with cathode-ray tubes?
49. What experimental results led to the conclusion that electrons were part of all forms of matter?



■ Figure 4.23


50. **Cathode Ray** Use the elements labeled in Figure 4.23 to explain the direction of a cathode ray inside a cathode-ray tube.
51. Briefly explain how Rutherford discovered the nucleus.
52. **Particle Deflection** What caused the deflection of the alpha particles in Rutherford's gold foil experiment?
53. **Charge of Cathode Rays** How was an electric field used to determine the charge of a cathode ray?
54. Explain what keeps the electrons confined in the space surrounding the nucleus.
55. What is the approximate size of an atom?
56. **Visualizing Atoms** What technique can be used to visualize individual atoms?
57. What are the strengths and weaknesses of Rutherford's nuclear model of the atom?

## Section 4.3

## Mastering Concepts

58. How do isotopes of a given element differ? How are they similar?
59. How is an atom's atomic number related to its number of protons? To its number of electrons?
60. How is the mass number related to the number of protons and neutrons an atom has?

61. How can you determine the number of neutrons in an atom if its mass number and its atomic number are known.
62. What do the superscript and subscript in the notation  ${}^{40}_{19}\text{K}$  represent?
63. **Standard Units** Define the atomic mass unit. What were the benefits of developing the atomic mass unit as a standard unit of mass?
64. **Isotopes** Are the following elements isotopes of each other? Explain.



65. Does the existence of isotopes contradict part of Dalton's original atomic theory? Explain.

### Mastering Problems

66. How many protons and electrons are contained in an atom of element 44?
67. **Carbon** A carbon atom has a mass number of 12 and an atomic number of 6. How many neutrons does it have?
68. **Mercury** An isotope of mercury has 80 protons and 120 neutrons. What is the mass number of this isotope?
69. **Xenon** An isotope of xenon has an atomic number of 54 and contains 77 neutrons. What is the xenon isotope's mass number?
70. If an atom has 18 electrons, how many protons does it have?
71. **Sulfur** Show that the atomic mass of the element sulfur is 32.065 amu.
72. Fill in the blanks in Table 4.6.

Element	Cl	Cl	Zr	Zr
Atomic number	17		40	
Mass number	35	37		92
Protons				40
Neutrons			50	
Electrons		17		

73. How many electrons, protons, and neutrons are contained in each atom?
- a.  ${}^{132}_{55}\text{Cs}$                       c.  ${}^{163}_{69}\text{Tm}$
- b.  ${}^{59}_{27}\text{Co}$                          d.  ${}^{70}_{30}\text{Zn}$
74. How many electrons, protons, and neutrons are contained in each atom?
- a. gallium-69                      c. titanium-48
- b. fluorine-23                      d. tantalum-181

75. For each chemical symbol, determine the number of protons and electrons an atom of the element contains.
- a. V                                      c. Ir
- b. Mn                                    d. S

76. Gallium, which has an atomic mass of 69.723 amu, has two naturally occurring isotopes, Ga-69 and Ga-71. Which isotope occurs in greater abundance? Explain.

77. **Atomic Mass of Silver** Silver has two isotopes:  ${}^{107}_{47}\text{Ag}$ , which has a mass of 106.905 amu and a percent abundance of 52.00%, and  ${}^{109}_{47}\text{Ag}$ , which has a mass of 108.905 amu and a percent abundance of 48.00%. What is the atomic mass of silver?

78. Data for chromium's four naturally occurring isotopes are provided in Table 4.7. Calculate chromium's atomic mass.

Table 4.7 Chromium Isotope Data

Isotope	Percent Abundance	Mass (amu)
Cr-50	4.35	49.946
Cr-52	83.79	51.941
Cr-53	9.50	52.941
Cr-54	2.36	53.939

## Section 4.4

### Mastering Concepts

79. What is radioactive decay?
80. Why are some atoms radioactive?
81. Discuss how radioactive atoms gain stability.
82. Define *alpha particle*, *beta particle*, and *gamma ray*.
83. Write the symbols used to denote alpha, beta, and gamma radiation and give their mass and charge.
84. What type of reaction involves changes in the nucleus of an atom?
85. **Radioactive Emissions** What change in mass number occurs when a radioactive atom emits an alpha particle? A beta particle? A gamma particle?
86. What is the primary factor that determines whether an nucleus is stable or unstable?
87. Explain how energy loss and nuclear stability are related to radioactive decay.
88. Explain what must occur before a radioactive atom stops to undergo further radioactive decay.
89. **Boron-10** emits alpha particles and cesium-137 emits beta particles. Write balanced nuclear reactions for each radioactive decay.

**Mixed Review**

- 90.** Determine what was wrong with Dalton's theory and provide the most recent version of the atomic structure.
- 91. Cathode-Ray Tube** Describe a cathode-ray tube and how it operates.
- 92. Subatomic Particles** Explain how J. J. Thomson's determination of the charge-to-mass ratio of the electron led to the conclusion that atoms were composed of subatomic particles.
- 93. Gold Foil Experiment** How did the actual results of Rutherford's gold foil experiment differ from the results he expected?
- 94.** If a nucleus contains 12 protons, how many electrons are in the neutral atom? Explain.
- 95.** An atom's nucleus has 92 protons and its mass number is 235. How many neutrons are in the nucleus? What is the name of the atom?
- 96.** Complete **Table 4.8**.

**Table 4.8** Composition of Various Isotopes

Isotope		Zn-64	
Atomic number		9	11
Mass number	32		23
Number of protons	16		
Number of neutrons		24	10
Number of electrons		20	


- 97.** Approximately how many times greater is the diameter of an atom than the diameter of its nucleus? Knowing that most of an atom's mass is contained in the nucleus, what can you conclude about the density of the nucleus?
- 98.** Is the charge of a nucleus positive, negative, or zero? The charge of an atom?
- 99.** Why are electrons in a cathode-ray tube deflected by electric fields?
- 100.** What was Henry Moseley's contribution to the modern understanding of the atom?
- 101.** What is the mass number of potassium-39? What is the isotope's charge?
- 102.** Boron-10 and boron-11 are the naturally occurring isotopes of elemental boron. If boron has an atomic mass of 10.81 amu, which isotope occurs in greater abundance?
- 103. Semiconductors** Silicon is important to the semiconductor manufacturing industry. The three naturally occurring isotopes of silicon are silicon-28, silicon-29, and silicon-30. Write the symbol for each.

- 104. Titanium** Use **Table 4.9** to calculate the atomic mass of titanium.

**Table 4.9** Titanium Isotopes

Isotope	Atomic Mass (amu)	Relative Abundance (%)
Ti-46	45.953	8.00
Ti-47	46.952	7.30
Ti-48	47.948	73.80
Ti-49	48.948	5.50
Ti-50	49.945	5.40

- 105.** Describe how each type of radiation affects an atom's atomic number and mass number.
- 106. Relative Abundances** Magnesium constitutes about 2% of Earth's crust and has three naturally occurring isotopes. Suppose you analyze a mineral and determine that it contains the three isotopes in the following proportions: Mg-24 (abundance = 79%), Mg-25 (abundance = 10%), and Mg-26 (abundance = 11%). If your friend analyzes a different mineral containing magnesium, do you expect her to obtain the same relative abundances for each magnesium isotope? Explain your reasoning.

**Figure 4.24**

- 107. Radiation** Identify the two types of radiation shown in **Figure 4.24**. Explain your reasoning.

**Think Critically**

- 108. Formulate** How were scientific methods used to determine the model of the atom? Why is the model considered a theory?
- 109. Discuss** What experiment led to the dispute of J. J. Thomson's plum pudding atomic model? Explain your answer.
- 110. Apply** Which is greater, the number of compounds or the number of elements? The number of elements or the number of isotopes? Explain.

- 111. Analyze** An element has three naturally occurring isotopes. What other information must you know in order to calculate the element's atomic mass?
- 112. Apply** If atoms are primarily composed of empty space, explain why you cannot pass your hand through a solid object.
- 113. Formulate** Sketch a modern atomic model of a typical atom and identify where each type of subatomic particle would be located.
- 114. Apply** Indium has two naturally occurring isotopes and an atomic mass of 114.818 amu. In-113 has a mass of 112.904 amu and an abundance of 4.3%. What is the identity and percent abundance of indium's other isotope?
- 115. Infer** Sulfur's average atomic mass is close to the whole number 32. Chlorine's average atomic mass is 35.453, which is not a whole number. Suggest a possible reason for this difference.

### Challenge Problem

- 116. Magnesium Isotopes** Compute the mass number,  $X$ , of the third isotope of magnesium given that the respective abundances of the naturally occurring isotopes are: 79.0%, 10%, and 11% for  ${}_{12}^{24}\text{Mg}$ ,  ${}_{12}^{25}\text{Mg}$ ,  ${}_{12}^X\text{Mg}$ . The relative atomic mass of magnesium is 24.305 amu.

### Cumulative Review

- 117.** How is a qualitative observation different from a quantitative observation? Give an example of each. (*Chapter 1*)
- 118.** A 1.0-cm<sup>3</sup> block of gold can be flattened to a thin sheet that averages  $3.0 \times 10^{-8}$  cm thick. What is the area (in cm<sup>2</sup>) of the flattened gold sheet? (*Chapter 2*)
- 119.** A piece of paper has an area of 603 cm<sup>2</sup>. How many sheets of paper would the sheet of gold mentioned in problem 118 cover? (*Chapter 2*)
- 120.** Classify each mixture as heterogeneous or homogeneous. (*Chapter 3*)
- salt water
  - vegetable soup
  - 14-K gold
  - concrete
- 121.** Determine whether each change is physical or chemical. (*Chapter 3*)
- Water boils.
  - A match burns.
  - Sugar dissolves in water.
  - Sodium reacts with water.
  - Ice cream melts.

### Additional Assessment

- 122. Television and Computer Screens** Describe how cathode rays are used to generate television and computer monitor images.
- 123. The Standard Model** The standard model of particle physics describes all of the known building blocks of matter. Research the particles included in the standard model. Write a short report describing the known particles and those thought to exist but not yet detected experimentally.
- 124. STM** Individual atoms can be seen using a sophisticated device known as a scanning tunneling microscope. Write a short report on how the scanning tunneling microscope works and create a gallery of this microscope's images from sources such as books, magazines, and the Internet.

### DBQ Document-Based Questions

**Zirconium** is a lustrous, gray-white metal. Because of its high resistance to corrosion and its low cross section for neutron absorption, it is often used in nuclear reactors. It can also be processed to produce gems that look like diamonds and are used in jewelry.

Table 4.10 shows the relative abundances of zirconium isotopes.

**Table 4.10** Relative Abundances of Zirconium Isotopes

Element	Relative Abundance
Zirconium-90	51.4
Zirconium-91	11.2
Zirconium-92	17.2
Zirconium-94	17.4
Zirconium-96	2.8

Data obtained from: Lide, David R., ed. 2005. *CRC Handbook of Chemistry and Physics*. Boca Raton: CRC Press.

- 125.** What is the mass number of each zirconium isotope?
- 126.** Compute the number of protons and neutrons for each zirconium isotope.
- 127.** Does the number of protons or neutrons remain the same for all isotopes? Explain.
- 128.** Based on the relative abundances of each isotope, predict to which isotope's mass the average atomic mass of zirconium is going to be closest.
- 129.** Calculate the weighted average atomic mass of zirconium.




## Standardized Test Practice

## Multiple Choice

- Which describes an atom of plutonium?
  - It can be divided into smaller particles that retain all the properties of plutonium.
  - It cannot be divided into smaller particles that retain all the properties of plutonium.
  - It does not possess all the properties of a larger quantity of plutonium.
  - It has an atomic number of 244.
- Neptunium's only naturally occurring isotope,  $^{237}_{93}\text{Np}$ , decays by emitting one alpha particle, one beta particle, and one gamma ray. What is the new atom formed from this decay?
  - $^{233}_{92}\text{U}$
  - $^{241}_{93}\text{Np}$
  - $^{233}_{90}\text{Th}$
  - $^{241}_{92}\text{U}$
- Which type of matter has a definite composition throughout and is made of more than one type of element?
  - heterogeneous mixture
  - homogeneous mixture
  - element
  - compound

Use the diagram below to answer Question 4.




Key	
	= Atom of Element A
	= Atom of Element B

- Which diagram shows a mixture?
  - X
  - Y
  - Z
  - both X and Z
- The Moon is approximately 384,400 km from Earth. What is this value in scientific notation?
  - $384.4 \times 10^3$  km
  - $3.844 \times 10^5$  km
  - $3.844 \times 10^{-5}$  km
  - $3844 \times 10^{-2}$  km

- Why does an atom have no net electric charge?
  - Its subatomic particles carry no electric charges.
  - The positively charged protons cancel out the negatively charged neutrons.
  - The positively charged neutrons cancel out the negatively charged electrons.
  - The positively charged protons cancel out the negatively charged electrons.
- How many neutrons, protons, and electrons does  $^{126}_{52}\text{Te}$  have?
  - 126 neutrons, 52 protons, and 52 electrons
  - 74 neutrons, 52 protons, and 52 electrons
  - 52 neutrons, 74 protons, and 74 electrons
  - 52 neutrons, 126 protons, and 126 electrons

Use the figure below to answer Question 7.



- Record the length of this paper clip to the appropriate number of significant digits.
  - 31 mm
  - 31.1 mm
  - 30.1 mm
  - 31.15 mm
- Element X has an unstable nucleus due to an overabundance of neutrons. All are likely to occur EXCEPT
  - element X will undergo radioactive decay.
  - element X will eventually become a stable, non-radioactive element.
  - element X will gain more protons to balance the neutrons it possesses.
  - element X will spontaneously lose energy.
- What makes up most of the volume of an atom?
  - protons
  - neutrons
  - electrons
  - empty space

## Short Answer

11. A 36.41-g sample of calcium carbonate ( $\text{CaCO}_3$ ) contains 14.58 g of calcium and 4.36 g of carbon. What is the mass of oxygen contained in the sample? What is the percent by mass of each element in this compound?

Use the table below to answer Questions 12 and 13.

Characteristics of Naturally Occurring Neon Isotopes			
Isotope	Atomic Number	Mass (amu)	Percent Abundance
$^{20}\text{Ne}$	10	19.992	90.48
$^{21}\text{Ne}$	10	20.994	0.27
$^{22}\text{Ne}$	10	21.991	9.25

12. For each isotope listed above, write the number of protons, electrons, and neutrons it contains.
13. Using the data in the table above, calculate the average atomic mass of neon.

## Extended Response


14. Assume that Element Q has the following three isotopes:  $^{248}\text{Q}$ ,  $^{252}\text{Q}$ , and  $^{259}\text{Q}$ . If the atomic mass of Q is 258.63, which of its isotopes is most abundant? Explain your answer.
15. Iodine-131 undergoes radioactive decay to form an isotope with 54 protons and 77 neutrons. What type of decay occurs in this isotope? Explain how you can tell.
16. You are given an aluminum cube. Your measurements show that its sides are 2.14 cm and its mass is 25.1 g. Explain how you would find its density. If the density of aluminum is known to be  $2.70 \text{ g/cm}^3$ , what is your percent error?

### NEED EXTRA HELP?

If You Missed Question . . .	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Review Section . . .	4.2	4.4	3.4	4.2	4.3	3.1	2.2	4.4	2.2	4.2	3.4	4.3	4.3	4.3	4.4	2.3	4.3	4.2	4.2	4.1	3.3

## SAT Subject Test: Chemistry

For each question below, indicate whether Statement I is true or false and indicate whether Statement II is true or false. If Statement II is a *correct explanation* of Statement I, write CE on your paper.



### Statement I

### Statement II

17. The two atoms of boron pictured above are isotopes	BECAUSE	they have the same number of protons but a different number of neutrons.
18. Most alpha particles shot at a piece of gold foil travel through it	BECAUSE	an atom has a large nucleus compared to its overall size.
19. A beam of neutrons is attracted to the charged plates surrounding it	BECAUSE	neutrons have no charge.
20. Carbon and oxygen can form either $\text{CO}$ or $\text{CO}_2$	BECAUSE	carbon and oxygen obey the law of definite composition.
21. A mixture of sand and water is heterogeneous	BECAUSE	water is a compound formed from hydrogen and oxygen.